

Reactions to Traumatic Events

At some time in your life, you or someone you know may experience a traumatic event. It is common to feel emotional aftershocks after a traumatic event. Sometimes aftershocks appear immediately, but they also can appear a few hours or even days later. In some cases, weeks or months may pass before stress reactions occur.

The signs and symptoms of a stress reaction may last days and occasionally months, depending on the severity of the traumatic event.

With understanding and support from family, friends and co-workers, stress reactions usually pass quickly. Occasionally, stress reactions can be so severe that professional assistance from a counselor may be needed. Below are some signs and symptoms of stress reactions.

PHYSICAL MENTAL

Fatigue Tendency to blame others

Twitches Confusion

Difficulty breathing Inability to make decisions

Elevated blood pressure Heightened or lowered alertness

Rapid heartbeat Poor concentration

Chest pain **Forgetfulness**

Headaches Trouble identifying known objects or

people Nausea/vomiting

Increase or decreased awareness of Thirst

surrounding Hunger

Poor problem-solving **Dizziness**

Loss of a sense of time, place or Excessive sweating

person

Chills Disturbed thinking

Fainting Nightmares



Contact Us

Your program provides access to tools and other resources online or call us directly to speak to a professional counselor who is available everyday and at any time to provide confidential assistance at no cost to you.

Visit MagellanHealth.com/member

EMOTIONAL

Anxiety

Grief

Denial

Survivor guilt/self blame

Emotional numbness

Uncertainty

Loss of emotional control

Fear of loss

Intense anger

Irritability

Helplessness

Mistrust

Feelings of worthlessness

Apathy/boredom

BEHAVIORAL

Changes in normal activities

Withdrawal from others

Emotional outbursts

Change in communication

Suspiciousness

Inability to rest

Substance abuse

Intensified startle reflex

Decreased personal hygiene

Appetite disturbance

Accident proneness

Don't ignore these symptoms. If gone too long unnoticed there could be more harm done and depression, anxiety or other disorders may appear.